



A Semantic Completeness Proof for Tableaux Modulo

R. Bonichon and O. Hermant
Université Pierre et Marie Curie Paris VI

LPAR '06, November 14, 2006

Outline

- 1 Deduction modulo
- 2 Tableaux modulo
 - Rules
 - Soundness
 - Systematic generation
- 3 Semantic Completeness
 - Semi- and partial valuations
 - Conditions for model construction
- 4 Conclusion
 - Cut elimination
 - Up and coming



Proving: Computation vs. Deduction

Computation

Deterministic search for a proof.

Deduction

Non-deterministic search for a proof.

Deduction modulo = computation + deduction

Deduction

Natural deduction, *sequent calculus*

Computation

- Term rewriting

$$S(x) + y \rightarrow S(x + y)$$

- Equational axioms

$$x + y = y + x$$

- **Propositional rewriting**

$$x * y = 0 \rightarrow x = 0 \vee y = 0$$

Extension of the sequent calculus

\mathcal{R} : set of propositional rewrite rules

\mathcal{E} : set of term rewrite rules and equational axioms.

$\equiv_{\mathcal{R}\mathcal{E}}$: congruence defined by $\mathcal{R} \cup \mathcal{E}$.

Sequent modulo rule

$$\frac{\Gamma, A \vdash_{\mathcal{R}\mathcal{E}} \Delta \quad \Gamma, B \vdash_{\mathcal{R}\mathcal{E}} \Delta}{\Gamma, P \vdash_{\mathcal{R}\mathcal{E}} \Delta} \vee\text{-I}$$

if $P \equiv_{\mathcal{R}\mathcal{E}} A \vee B$.

Benefits

- Shorter more readable proofs

$\overline{\Gamma, 4 = 4 \vdash_{\mathcal{RE}} 2 + 2 = 4, \Delta}$ axiom, $\mathcal{RE} \hat{=} \text{Peano's arithmetic}$

- Powerful general framework for theories without axioms
 - HOL (Dowek, Hardin, Kirchner, MSCS 01)
 - (Heyting's) arithmetic (Dowek, Werner, RTA '05)
 - Zermelo's set theory (Dowek, Miquel)



Tableaux modulo: what we want

- Better free variable tableaux for deduction modulo (Bonichon, IJCAR'04)
- Term rewriting and propositional rewriting in \mathcal{R}
- Soundness (semantic)
- Completeness (semantic)



Expanding tableaux

- Free-variable constrained tableaux of constrained formulas
- Rules for:
 - Connectors: α and β rules
 - Quantifiers: γ and δ rules
 - Rewriting (on terms and propositions)
 - Closure



Rules

$$\frac{\alpha(A, B)}{A, B}$$

$$\frac{\alpha(A, B)}{A \wedge B}$$

$$\neg(A \vee B)$$

$$\neg(A \Rightarrow B)$$



Rules

$$\frac{\alpha(A, B)}{A, B}$$

$$\frac{\beta(A, B)}{A \mid B}$$

$$\frac{\beta(A, B)}{A \vee B}$$

$$\neg(A \wedge B)$$

$$A \Rightarrow B$$



Rules

$$\frac{\alpha(A, B)}{A, B}$$

$$\frac{\beta(A, B)}{A \mid B}$$

$$\frac{\gamma(x, A)}{\forall x A}$$

$$\neg(\exists x A)$$

$$\frac{\gamma(x, A)}{A[x := X]}$$



Rules

$$\frac{\alpha(A, B)}{A, B}$$

$$\frac{\beta(A, B)}{A \mid B}$$

$$\frac{\delta(x, A)}{\exists x A}$$

$$\neg(\forall x A)$$

$$\frac{\gamma(x, A)}{A[x := X]}$$

$$\frac{\delta(x, A)}{A[x := f_{\text{sko}}(\text{args})]}$$



Rewriting

$\mathcal{T} \cdot \mathcal{C}$: tableau \mathcal{T} with global constraint \mathcal{C}

P_c : proposition P with local constraint c

$$\frac{\Gamma_1, P_c \mid \dots \mid \Gamma_n \cdot \mathcal{C}}{\Gamma_1, P_c, P_{c'}[r]_\omega \mid \dots \mid \Gamma_n \cdot \mathcal{C}} \text{rw}$$

if $l \rightarrow r$, and $c' = c \cup \{P|_\omega \stackrel{?}{=} l\}$



Soundness

Theorem (Soundness)

If Γ has a \mathcal{R} -model, then we can not derive the closed tableau \odot from Γ using the rules for tableaux modulo.

Proof.

As usual. \mathcal{R} -models are extensions of boolean models such that for any propositions $P \equiv_{\mathcal{R}} Q$, $|P| = |Q|$. □



Systematic tableau expansion

$$\mathcal{R} = P(a) \rightarrow Q(a)$$

Rule:

$$\forall x P(x) \wedge (\neg P(c) \vee \neg Q(a))$$

$$\mathcal{C} = \emptyset$$



Systematic tableau expansion

$$\forall x P(x) \wedge (\neg P(c) \vee \neg Q(a))$$

$$\forall x P(x)$$

$$\neg P(c) \vee \neg Q(a)$$

$$\mathcal{R} = P(a) \rightarrow Q(a)$$

$$\text{Rule: } \frac{\alpha(A, B)}{A, B}$$

$$\mathcal{C} = \emptyset$$



Systematic tableau expansion

$$\forall x P(x) \wedge (\neg P(c) \vee \neg Q(a))$$

$$\begin{array}{c}
 \forall x P(x) \\
 | \\
 \neg P(c) \vee \neg Q(a) \\
 \swarrow \quad \searrow \\
 \neg P(c) \quad \neg Q(a)
 \end{array}$$

$$\mathcal{R} = P(a) \rightarrow Q(a)$$

$$\text{Rule: } \frac{\beta(A, B)}{A \mid B}$$


$$\mathcal{C} = \emptyset$$

	l	r
lgth	4	4
$\gamma > \text{rw}$	1	1



Systematic tableau expansion

$$\forall x P(x) \wedge (\neg P(c) \vee \neg Q(a))$$



$$\mathcal{R} = P(a) \rightarrow Q(a)$$

$$\text{Rule: } \frac{\gamma(x, A)}{A[x := X]}$$


$$\mathcal{C} = \emptyset$$

	l	r
lgth	5	4
$\gamma > \text{rw}$	0	1



Systematic tableau expansion

$$\forall x P(x) \wedge (\neg P(c) \vee \neg Q(a))$$



$$\mathcal{R} = P(a) \rightarrow Q(a)$$

$$\text{Rule: } \frac{\gamma(x, A)}{A[x := X]}$$


$$\mathcal{C} = \emptyset$$

	l	r
lgth	5	5
$\gamma > \text{rw}$	0	0



Systematic tableau expansion

$$\forall x P(x) \wedge (\neg P(c) \vee \neg Q(a))$$



$$\mathcal{R} = P(a) \rightarrow Q(a)$$

$$\text{Rule: } \frac{P(l)}{P(r)}$$


$$\mathcal{C} = \emptyset$$

	l	r
lgth	6	5
$\gamma > \text{rw}$	1	0



Systematic tableau expansion

$$\forall x P(x) \wedge (\neg P(c) \vee \neg Q(a))$$



$$\mathcal{R} = P(a) \rightarrow Q(a)$$

$$\text{Rule: } \frac{P(l)}{P(r)}$$


$$\mathcal{C} = \emptyset$$

	l	r
lgth	6	6
$\gamma > rw$	1	1



Systematic tableau expansion

$$\forall x P(x) \wedge (\neg P(c) \vee \neg Q(a))$$



$$\mathcal{R} = P(a) \rightarrow Q(a)$$

Rule: **closure**

$$\mathcal{C} = \{X \stackrel{?}{=} c, Y \stackrel{?}{=} a\}$$

	l	r
lgth	6	6
$\gamma > rw$	1	1



Proving completeness

Theorem (Completeness)

Let \mathcal{R} be a confluent terminating rewrite system, and Γ a set of propositions. If the systematic tableau procedure rooted at Γ has an open branch, then Γ has a \mathcal{R} -model.



Usual model construction

From a complete open branch:

- Semi-valuation (Hintikka set)
- Model

And that's it !



Is that enough?

- Models must be models of the rewrite rules too!
- The intended meaning of $P \rightarrow Q$ is $P \Leftrightarrow Q$
- But we begin by defining the usual semi-valuation V from the open branch ...

Some problems

\mathcal{R} -semi-valuation (\mathcal{V})

When $P \rightarrow Q$ (using $l \rightarrow r$), $\mathcal{V}(P)$ must be defined when $\mathcal{V}(Q)$ is.

\mathcal{R} -partial valuation ($\tilde{\mathcal{V}}$)

We would like to have $\mathcal{V}(\forall x(A(x) \wedge B(x))) = 1$ when for every term t we have $\mathcal{V}(A(t)) = 1$ and $\mathcal{V}(B(t)) = 1$.

Confluence and termination of \mathcal{R}

Not enough for completeness (Dowek-Werner, Hermant) !



Positivity

Condition

Propositional rewrite rules $l \rightarrow r \in \mathcal{R}$ are such that all atoms occurring in r occur positively

Model construction

- if A is an atom, and if $\tilde{V}(A)$ is defined, set $|A|_{\mathcal{R}} = \tilde{V}(A)$.
- if A is an atom, and if $\tilde{V}(A)$ is not defined, set $|A|_{\mathcal{R}} = 1$.
- if P is a compound proposition, set $|P|_{\mathcal{R}}$ accordingly.



Order

Condition

We consider a confluent rewrite system and a well-founded order \prec such that:

- if $P \rightarrow Q$ then $Q \prec P$.
- if A is a subformula of B then $A \prec B$.

Model construction

- if A is a normal atom, set $|A|_{\mathcal{R}} = \tilde{V}(A)$, if defined. Else set $|A|_{\mathcal{R}}$ arbitrarily.
- if A is not a normal atom, set $|A|_{\mathcal{R}} = |A \downarrow|_{\mathcal{R}}$.
- if P is a compound proposition, set $|P|_{\mathcal{R}}$ from the interpretation of its immediate subformulas.





Mixed

Condition

Consider two rewrite systems $\mathcal{R}_>$ and \mathcal{R}_+ .

$\mathcal{R} = \mathcal{R}_> \cup \mathcal{R}_+$ must be confluent and terminating.

\mathcal{R}_+ is right-normal for $\mathcal{R}_>$

Right-normality

Let two rewrite systems \mathcal{R}' and \mathcal{R} . \mathcal{R}' is right normal for \mathcal{R} if, for any propositional rule $l \rightarrow r \in \mathcal{R}'$, all the instances of atoms of r by \mathcal{R} -normal substitutions σ are in normal form for \mathcal{R} .



What we have

- Correct and complete tableau method for deduction modulo
- Somewhat better than the previous one
- Extract classes of rewrite systems



Cut elimination


- Not always normalization

$$R \in R \longrightarrow \forall y (\forall x (y \in x \Rightarrow R \in x) \Rightarrow (y \in R \Rightarrow (A \Rightarrow A)))$$

- Semantic cut elimination (admissibility)




Cut elimination







Cut elimination



Cut elimination



Cut elimination





Further research

- Companion paper on intuitionistic logic: constructive cut elimination
- More classes for semantic completeness
- Using semantic methods for HOL-C (with combinators)
- Ongoing implementation in Zenon prover (Doligez)